

```

!*** Two arrays must be declared:
!*** y (2) - an array containing the coordinate x = y (1) and speed V = y (2) and the working array
!*** work (15) with a length to be calculated using the equation  $3 + 6 * neqn$  (neqn- is number of
!*** equations)
!*** mu_atm and m_keha variables must also be declared as REAL. Here we take into
!*** account the Fortran feature. If variables are not declared at the beginning of the program by the
!*** REAL, INTEGER, etc. operators, the fortran compiler can declare and create them by using the
!*** default rule: if the first character in the variable name is i, j, k, l, m, n then the variable must be
!*** INTEGER if it is not so then the variable must be REAL
real y(2),work(15),mu_atm,m_keha
!*** Declaration of an additional integer array iwork(5) with fixed length 5
integer iwork(5)
!*** Description the global variables which must be used in subroutine "func" to calculate
!*** derivatives of coordinate x and velocity v with respect to time
common g_rask,ro_atm_0,ro_keha,param1,param2
!*** Next operator should be used to declare that "func" is an external subroutine,
!*** not a regular variable name
external func
!*** Now we need to open the file to save the calculations results
open(10,file="lang.dat")
!*** On the next lines some important variables must be created and we must assign
!*** values for them
!*** value of pi
pi=3.1415926536
!*** Acceleration of gravity for Tallinn
g_rask=9.814
!*** Radius of sphere (object of investigation) in meters
r_keha=1.8
!*** Sphere square of cross section
s_keha=pi*r_keha**2
!*** Aerodynamic parameter for sphere (taken from the internet)
c_keha=0.47
!*** volume of the sphere
v_keha=4.*pi*r_keha**3/3
!*** Mass of the sphere in kilograms
m_keha=15.
!*** Density of sphere
ro_keha=m_keha/v_keha
!*** Pressure of atmosphere on the sea level in Pascale
p_atm=100000.
!*** Temperature of atmosphere on the sea level in Kelvin
t_atm=293.
!*** Mass of one kilomole of air (average value) in kg/kmol
mu_atm=29.

```

```

**** Universal gas constant in  $\frac{J}{K \cdot kmol}$ 
r_gaas=8314.
**** Calculation of the density of air on the sea level
ro_atm_0=p_atm*mu_atm/(r_gaas*t_atm)
**** Two additional work variables used for convenience only
param1=g_rask*mu_atm/(r_gaas*t_atm)
param2=0.5*c_keha*s_keha/m_keha
**** Number of differential equations to be integrated
neqn=2
**** Total number of time steps
nt=400
**** Time step in seconds
dt=0.1
**** Initial value for time
t=0.
**** Relative and absolute errors for calculation the coordinate and velocity
relerr=1.e-7
abserr=1.e-7
**** Initial conditions
**** y(1)=1 - position of sphere at start time moment
**** y(2)=0 - velocity of sphere at start time moment
y(1)=1.
y(2)=0.
**** iflag=1 is need to start the calculation, it means that we are starting a new simulation
iflag=1

**** Cycle operator to perform the integration of the system of differential equations and
**** calculation of coordinate and velocity at different time moments with timestep dt=0.1 seconds.
**** The total time of simulation can be calculated as a product of variables nt and dt variables,
**** in our case it is equal to nt*dt=0.1*400=40 seconds.
do it=1,nt
**** Caslcluation of new value of time (should be done by hand)
tout=t+dt
**** Calculation of coordinate y(1) and velocity y(2) at the next time moment tout=t+dt
call rkf45(func, neqn, y, t, tout, relerr, abserr, iflag, work, iwork)
**** Just in case the control of the variable iflag value. If iflag=2 then calculation was success and
**** we can continue
if (iflag.ne.2) then
iflag=2
endif
**** if-operator to control the position of ball. If coordinate (height of trajectory) is less
**** than zero we have to stop calculations. In our case the height and x-coordinate of
**** ball are coincided.

```

```

if(y(1).le.0.) then
stop
endif
!*** Now we need to save the calculated results at the next format of data
!*** time x V
!*** 3 real numbers on each line and total number of lines is equal to value of variable "nt"
write(10,*) tout,y
!*** End of cycle operator
enddo
!*** stop program
stop
!*** End of source code for main program
end

!*** The most important part of the program. Here we must implement the differential equation.
!*** Now we !*** need to calculate the derivatives with respect to the coordinate and velocity
!*** with respect to time.
!*** Now the subroutine "func" must be created. Number of input parameters is fixed.
!*** t-time, y(2)-array consist the coordinate y(1) and velocity y(2) of our sphere,
!*** yp(2)-array with derivatives so that:
!***  $yp(1)=dy(1)/dt=velocity=y(2)$  and
!***  $yp(2)=dy(2)/dt=acceleration=g\_rask*(roh/ro\_keha-1.)-param2*roh*v*y(2)$ 
!*** (you can find a theoretical justification in the lecture materials)
subroutine func(t,y,yp)
!*** Declaration of arrays
real y(2),yp(2)
!*** Globalization of some variables (same as in the main program)
common g_rask,ro_atm_0,ro_keha,param1,param2
!*** first derivative for coordinate x
yp(1)=y(2)
!*** Calculation of the density of air on height y(1)
roh=ro_atm_0*exp(-param1*y(1))
!*** Absolute value of velocity
v=abs(y(2))
!*** Second derivative, calculation of acceleration with the second Newton's law
yp(2)=g_rask*(roh/ro_keha-1.)-param2*roh*v*y(2)
!*** Back to call operator in the main program
return
!*** End of source code for subroutine func
end

```